

SAGA APS

STUDY REPORT 2020-9159/20 23 00925

Surface Sanitizer Formula 80%

SUSPENSION TEST ACCORDING TO EN 1276:2019 (Phase 2 step 1)

Chemical disinfectants and antiseptics
Quantitative suspension test for the evaluation of bactericidal activity of chemical disinfectants and antiseptics used in food, industrial, domestic and institutional areas – Test method and requirements (phase 2, step 1)

NOVEMBER 2020

QACS LTD, 1 Antigonis str, 144-51 Metamorfossi, Greece. Tel: +30-210-2934745,
fax: +30-210-2934606, email: info@qacs.gr website: www.qacslab.com

STUDY REPORT 2020-9159/20 23 00925

SUSPENSION TEST ACCORDING TO EN 1276:2019

Chemical disinfectants and antiseptics - Quantitative suspension test for the evaluation of bactericidal activity of chemical disinfectants and antiseptics used in food, industrial, domestic and institutional areas - Test method and requirements (phase 2, step 1)

TEST PRODUCT IDENTIFICATION

| | | |
|-------------------------------------|---|-------------------------------|
| PRODUCT NAME | : | Surface Sanitizer Formula 80% |
| SUBSTANCES AND THEIR CONCENTRATIONS | : | Ethanol 80% w/w |
| APPEARANCE OF THE PRODUCT | : | Liquid |
| STORAGE CONDITIONS | : | Room Temperature, Darkness |
| LOT | : | Not Provided |
| METHOD | : | EN 1276:2019 |
| CONTACT TIME | : | 30 seconds |
| CONCENTRATION | : | Undiluted (80%), 50%, 1%. |
| STUDY SPONSOR | : | Saga ApS |
| PRODUCT SUPPLIER | : | Saga ApS |
| PRODUCT MANUFACTURER | : | Saga ApS |
| RECEIPT DATE | : | 21/09/2020 |
| STUDY PERIOD | : | 13/10/2020-15/10/2020 |
| LAB ID | : | 2020-9159/20 23 00925 |

SCOPE

This document specifies a test method and the minimum requirements for bactericidal activity of chemical disinfectant and antiseptic products that form a homogeneous, physically stable preparation when diluted with hard water or - in the case of ready-to-use products - with water. Products can only be tested at a concentration of 80 % or less, as some dilution is always produced by adding the test organisms and interfering substance.

This document applies to products that are used in food, industrial, domestic and institutional areas excluding areas and situations where disinfection is medically indicated and excluding products used on living tissues except those for hand hygiene in the above considered areas

PRINCIPLE

A sample of the product as delivered and/or diluted with hard water (or water for ready-to-use products with the exception of handwash products whose first dilution is done in hard water is added to a test suspension of bacteria in a solution of an interfering substance. The mixture is maintained at the chosen test temperature for the adopted contact time. At the end of this contact time, an aliquot is taken, and the bactericidal and/or the bacteriostatic activity in this portion is immediately neutralized or suppressed by a validated method. The method of choice is dilution-neutralization. If a suitable neutralizer cannot be found, membrane filtration is used. The numbers of surviving bacteria in each sample are determined and the reduction is calculated.

The test is performed using *Pseudomonas aeruginosa*, *Escherichia coli*, *Staphylococcus aureus* and *Enterococcus hirae* as test organisms. For temperatures ≥ 40 °C only *Enterococcus faecium* shall be used. For testing of hand hygiene products, *Pseudomonas aeruginosa*, *Escherichia coli* K12, *Staphylococcus aureus* and *Enterococcus hirae* are used as test organisms.

TEST CONDITIONS

1. The following procedure was performed in water bath at 20 °C
2. The test product was tested at 30 seconds contact time
3. A final concentration of 0.3g/L bovine albumin was used for testing (clean conditions)
4. Neutralization Method used: Dilution neutralization.
5. Neutralizer used: LPT Dilution Broth containing polysorbate 80.
6. According to EN 1276, products shall be tested at a minimum of three different concentrations to include one concentration in the active range and one concentration in the non-active range. In this case the product was tested: Undiluted (80%), 50%, 1%.

TEST ORGANISMS

| | |
|-------------------------------|-------------|
| <i>Pseudomonas aeruginosa</i> | NCIMB 10421 |
| <i>Staphylococcus aureus</i> | ATCC 6538 |
| <i>Escherichia coli</i> | NCIMB 8879 |
| <i>Enterococcus hirae</i> | NCIMB 8192 |

BACTERICIDAL ACTIVITY FOR GENERAL PURPOSES

The product shall be deemed to have passed the EN 1276 standard if it demonstrates in a valid test at least a 5 lg reduction, under the suitable test conditions for general purpose defined by this standard when the test organisms are *Pseudomonas aeruginosa*, *Escherichia coli*, *Staphylococcus aureus* and *Enterococcus hirae* (*E. faecium* when the test temperature is ≥ 40 °C).

ASSAY ACCEPTANCE CRITERIA

1. Test Suspension (N) is between 1.5 to 5.0 X 10⁸ CFU per mL ($8.17 \leq \log N \leq 8.70$)
2. No (N/10) is between 1.5 to 5.0 X 10⁷ CFU per mL ($7.17 \leq \log No \leq 7.70$)
3. Validation Suspension=Nv is between 3.0 x 10² and 1.6 x 10³.
4. Nvo (Nv/10) is between 30 and 160
5. Na is the number of survivors (cells) per ml in the test mixture at the end of contact time.
6. R (log reduction) = No - Na
7. Average recovery values for the experimental conditions control (A) were equal to or greater than 0.5 times the Validation Suspension (Nvo)
8. Average recovery values for the Neutralizer control (B) were equal to or greater than 0.5 times the Validation Suspension (Nvo)
9. Average recovery values for the Method Validation control (C) were equal to or greater than 0.5 times the Validation Suspension (Nvo)
10. Control of weighted mean counts. Quotient is not lower than 5 and not higher than 15

ARCHIVING

The laboratory book which contains all the information (raw data and results) regarding the study and the study reports are kept in the laboratory archives for 5 years.

TEST RESULTS FOR *Pseudomonas aeruginosa* (BACTERICIDAL SUSPENSION TEST)

Test suspension

| Test - suspension (N and No) | | | | |
|------------------------------|-----|-----|-------------------------|----------|
| N | Vc1 | Vc2 | x mean | 2.30E+08 |
| 10 ⁻⁶ | 237 | 222 | | |
| 10 ⁻⁷ | 22 | 24 | log N | 8.36 |
| | | | No (N/10) | 2.30E+07 |
| | | | log No | 7.36 |
| | | | 7,17 < = logNo < = 7,70 | Yes |

Validation and controls

| Validation suspension (Nvo) | | Experimental conditions (A) | | Neutralizer control (B) | | Method validation (C) Undiluted Product conc.: (80%) | |
|-----------------------------|--------|------------------------------------|--------|--|--------|--|--------|
| VC 1 | 48 | VC 1 | 59 | VC 1 | 48 | VC 1 | 52 |
| | x mean | | x mean | | x mean | | x mean |
| VC 2 | 59 | VC 2 | 42 | VC 2 | 45 | VC 2 | 65 |
| | 53.5 | | 50.5 | | 46.5 | | 58.5 |
| 30<x mean of Nvo < 160? | | x mean of A is >0,5*x mean of Nvo? | | x mean of B is >0,5*x mean of Nvo or Nvo/1000? | | x mean of C is >0,5*x mean of Nvo? | |
| Yes | | Yes | | Yes | | Yes | |

Test Results

| Product concentration (%) | Contact time | Dilution step | Vc 1 | Vc 2 | Average of Vc1 and Vc2 | Na= average x10 | log Na | log No | log Reduction (No-Na) | Criteria | Result |
|---------------------------|--------------|------------------|-------|-------|------------------------|-----------------|--------|--------|-----------------------|----------|------------|
| Undiluted (80%) | 30 seconds | 10 ⁻² | 0 | 0 | < 14 | < 140 | < 2.15 | 7.36 | > 5.21 | ≥ 5 | PASS TEST |
| | | 10 ⁻¹ | 0 | 0 | | | | | | | |
| 50% | 30 seconds | 10 ⁻² | 12 | 14 | 15 | 150 | 2.18 | 7.36 | 5.18 | ≥ 5 | PASS TEST |
| | | 10 ⁻¹ | 1 | 2 | | | | | | | |
| 1% | 30 seconds | 10 ⁻² | > 330 | > 330 | > 3300 | > 33000 | > 4.52 | 7.36 | < 2.84 | ≥ 5 | FAILS TEST |
| | | 10 ⁻¹ | > 330 | > 330 | | | | | | | |

TEST RESULTS FOR *Staphylococcus aureus* (BACTERICIDAL SUSPENSION TEST)

Test suspension

| Test - suspension (N and No) | | | |
|---------------------------------|-----|-----|-----------------------------|
| N | Vc1 | Vc2 | |
| 10^{-7} | 35 | 40 | x mean 3.77E+08 |
| 10^{-8} | 4 | 4 | log N 8.58 |
| | | | No (N/10) 3.77E+07 |
| | | | log No 7.58 |
| | | | 7,17 < = logNo < = 7,70 Yes |

Validation and controls

| Validation suspension (Nvo) | | Experimental conditions (A) | | Neutralizer control (B) | | Method validation (C) Undiluted Product conc.: (80%) | | |
|--------------------------------|-----|------------------------------------|----------|---|---------|---|---------|--------|
| VC 1 | 92 | x mean | VC 1 100 | x mean | VC 1 88 | x mean | VC 1 92 | x mean |
| VC 2 | 102 | 97 | VC 2 92 | 96 | VC 2 89 | 88.5 | VC 2 94 | 93 |
| 30<x mean of Nvo < 160? | | x mean of A is >0,5*x mean of Nvo? | | x mean of B is > 0,5*x mean of Nvo or Nvb/1000? | | x mean of C is >0,5*x mean of Nvo? | | |
| Yes | | Yes | | Yes | | Yes | | |

Test Results

| Product concentration (%) | Contact time | Dilution step | Vc 1 | Vc 2 | Average of Vc1 and Vc2 | Na= average x10 | log Na | log No | log Reduction (No-Na) | Criteria | Result |
|---------------------------|--------------|---------------|-------|-------|---------------------------|--------------------|--------|--------|-----------------------------|----------|------------|
| Undiluted (80%) | 30 seconds | 10^0 | 0 | 0 | < 14 | < 140 | < 2.15 | 7.58 | > 5.43 | ≥ 5 | PASS TEST |
| | | 10^{-1} | 0 | 0 | | | | | | | |
| 50% | 30 seconds | 10^0 | 0 | 0 | < 14 | < 140 | < 2.15 | 7.58 | > 5.43 | ≥ 5 | PASS TEST |
| | | 10^{-1} | 0 | 0 | | | | | | | |
| 1% | 30seconds | 10^0 | > 330 | > 330 | > 3300 | > 33000 | > 4.52 | 7.58 | < 3.06 | ≥ 5 | FAILS TEST |
| | | 10^{-1} | > 330 | > 330 | | | | | | | |

TEST RESULTS FOR *Escherichia Coli* (BACTERICIDAL SUSPENSION TEST)

Test suspension

| Test - suspension (N and No) | | | | |
|---------------------------------|-----|-----|-------------------------|----------|
| N | Vc1 | Vc2 | x mean | 4.64E+08 |
| 10 ⁻⁷ | 46 | 48 | | |
| 10 ⁻⁸ | 4 | 4 | log N | 8.67 |
| | | | No (N/10) | 4.64E+07 |
| | | | log No | 7.67 |
| | | | 7,17 < = logNo < = 7,70 | Yes |

Validation and controls

| Validation suspension (Nvo) | | | Experimental conditions (A) | | | Neutralizer control (B) | | | Method validation (C) Undiluted Product conc.: (80%) | | |
|--------------------------------|----|--------|-------------------------------------|-----|--------|---|-----|--------|---|----|--------|
| VC 1 | 92 | x mean | VC 1 | 98 | x mean | VC 1 | 97 | x mean | VC 1 | 96 | x mean |
| VC 2 | 99 | 95,5 | VC 2 | 102 | 100 | VC 2 | 104 | 100,5 | VC 2 | 99 | 97,5 |
| 30*x mean of Nvo < 160? | | | x mean of A is > 0,5*x mean of Nvo? | | | x mean of B is > 0,5*x mean of Nvo or Nva/1000? | | | x mean of C is > 0,5*x mean of Nvo? | | |
| Yes | | | Yes | | | Yes | | | Yes | | |

Test Results

| Product concentration (%) | Contact time | Dilution step | Vc 1 | Vc 2 | Average of Vc1 and Vc2 | Na= average x10 | log Na | log No | log Reduction (No-Na) | Criteria | Result |
|---------------------------|--------------|------------------|-------|-------|---------------------------|--------------------|--------|--------|-----------------------------|----------|------------|
| Undiluted (80%) | 30 seconds | 10 ⁰ | 0 | 0 | < 14 | < 140 | < 2.15 | 7.67 | > 5.52 | ≥ 5 | PASS TEST |
| | | 10 ⁻¹ | 0 | 0 | | | | | | | |
| 50% | 30 seconds | 10 ⁰ | 0 | 0 | < 14 | < 140 | < 2.15 | 7.67 | > 5.52 | ≥ 5 | PASS TEST |
| | | 10 ⁻¹ | 0 | 0 | | | | | | | |
| 1% | 30 seconds | 10 ⁰ | > 330 | > 330 | > 3300 | > 33000 | > 4.52 | 7.67 | < 3.15 | ≥ 5 | FAILS TEST |
| | | 10 ⁻¹ | > 330 | > 330 | | | | | | | |

TEST RESULTS FOR *Enterococcus hirae* (BACTERICIDAL SUSPENSION TEST)

Test suspension

| Test - suspension (N and No) | | | |
|------------------------------|-----|-----|-----------------------------|
| N | Vc1 | Vc2 | |
| 10 ⁻⁶ | 259 | 272 | x mean 2.64E+08 |
| 10 ⁻⁷ | 28 | 22 | log N 8.42 |
| | | | No (N/10) 2.64E+07 |
| | | | log No 7.42 |
| | | | 7,17 < = logNo < = 7,70 Yes |

Validation and controls

| Validation suspension (Nvo) | | | Experimental conditions (A) | | | Neutralizer control (B) | | | Method validation (C) Undiluted Product conc.: (80%) | | |
|-----------------------------|----|--------|-------------------------------------|----|--------|---|----|--------|--|----|--------|
| VC 1 | 48 | x mean | VC 1 | 59 | x mean | VC 1 | 48 | x mean | VC 1 | 52 | x mean |
| VC 2 | 59 | 53.5 | VC 2 | 42 | 50.5 | VC 2 | 45 | 46.5 | VC 2 | 65 | 58.5 |
| 30*x mean of Nvo < 160? | | | x mean of A is > 0,5*x mean of Nvo? | | | x mean of B is > 0,5*x mean of Nvo or Nva/1000? | | | x mean of C is > 0,5*x mean of Nvo? | | |
| Yes | | | Yes | | | Yes | | | Yes | | |

Test Results

| Product concentration (%) | Contact time | Dilution step | Vc 1 | Vc 2 | Average of Vc1 and Vc2 | Na= average x10 | log Na | log No | log Reduction (No-Na) | Criteria | Result |
|---------------------------|--------------|------------------|-------|-------|------------------------|-----------------|--------|--------|-----------------------|----------|------------|
| Undiluted (80%) | 30 seconds | 10 ⁻² | 0 | 0 | < 14 | < 140 | < 2.15 | 7.42 | > 5.28 | ≥ 5 | PASS TEST |
| | | 10 ⁻¹ | 0 | 0 | | | | | | | |
| 50% | 30 seconds | 10 ⁻² | 0 | 0 | < 14 | < 140 | < 2.15 | 7.42 | > 5.28 | ≥ 5 | PASS TEST |
| | | 10 ⁻¹ | 0 | 0 | | | | | | | |
| 1% | 30seconds | 10 ⁻² | > 330 | > 330 | > 3300 | > 33000 | > 4.52 | 7.42 | < 2.90 | ≥ 5 | FAILS TEST |
| | | 10 ⁻¹ | > 330 | > 330 | | | | | | | |

CONCLUSION

TEST SUBSTANCE IDENTIFICATION

| | |
|-------------------------------------|---------------------------------|
| PRODUCT NAME | : Surface Sanitizer Formula 80% |
| SUBSTANCES AND THEIR CONCENTRATIONS | : Ethanol 80% w/w |
| APPEARANCE OF THE PRODUCT | : Liquid |
| STORAGE CONDITIONS | : Room Temperature, Darkness |
| LOT | : Not Provided |
| METHOD | : EN 1276:2019 |
| CONTACT TIME | : 30 seconds |
| CONCENTRATION | : Undiluted (80%), 50%, 1%. |
| STUDY SPONSOR | : Saga ApS |
| PRODUCT SUPPLIER | : Saga ApS |
| PRODUCT MANUFACTURER | : Saga ApS |
| RECEIPT DATE | : 21/09/2020 |
| STUDY PERIOD | : 13/10/2020-15/10/2020 |
| LAB ID | : 2020-9159/20 23 00925 |

METHODOLOGY ABSTRACT


A test suspension of bacteria is tested against a product test solution at three different concentrations with the presence of interfering substance. The mixture is maintained at 20°C ±1°C for 30 seconds. At the end of this contact time, an aliquot is taken, and the bactericidal activity in this portion is immediately neutralized or suppressed. The numbers of surviving flora are determined and the log reduction is calculated.

RESULT

The product under test: "Surface Sanitizer Formula 80%" demonstrated bactericidal activity according to EN 1276:2019 (≥ 5 log reduction), under clean conditions for 30 seconds contact time, at 20 ± 1 °C, when tested at product concentrations:

Undiluted (80%) using as test organisms the reference strains: *Pseudomonas aeruginosa*, *Staphylococcus aureus*, *Escherichia coli* and *Enterococcus hirae*.

For the QACS Ltd Laboratory,


QACS Laboratories
1 Antigonis str, 144 51 Metamorfossi Greece
VAT no EL 998700411, email: info@qacs.gr
Tel +30-2102934745 fax +30-210 2934606
www.qacs.gr

Signature date: 09/11/2020
Lagiopoulos Giorgos
Agronomist-Food Technologist M.Sc.
Study Manager

STUDY SUMMARY / ABSTRACT

SUSPENSION TEST ACCORDING TO EN 1276:2019

Chemical disinfectants and antiseptics - Quantitative suspension test for the evaluation of bactericidal activity of chemical disinfectants and antiseptics used in food, industrial, domestic and institutional areas - Test method and requirements (phase 2, step 1)

| | | |
|-------------------------------------|---|-------------------------------|
| PRODUCT NAME | : | Surface Sanitizer Formula 80% |
| SUBSTANCES AND THEIR CONCENTRATIONS | : | Ethanol 80% w/w |
| APPEARANCE OF THE PRODUCT | : | Liquid |
| STORAGE CONDITIONS | : | Room Temperature, Darkness |
| LOT | : | Not Provided |
| METHOD | : | EN 1276:2019 |
| CONTACT TIME | : | 30 seconds |
| CONCENTRATION | : | Undiluted (80%), 50%, 1%. |
| STUDY SPONSOR | : | Saga ApS |
| PRODUCT SUPPLIER | : | Saga ApS |
| PRODUCT MANUFACTURER | : | Saga ApS |
| RECEIPT DATE | : | 21/09/2020 |
| STUDY PERIOD | : | 13/10/2020-15/10/2020 |
| LAB ID | : | 2020-9159/20 23 00925 |

TEST MICROORGANISMS

| | |
|-------------------------------|-------------|
| <i>Pseudomonas aeruginosa</i> | NCIMB 10421 |
| <i>Staphylococcus aureus</i> | ATCC 6538 |
| <i>Escherichia coli</i> | NCIMB 8879 |
| <i>Enterococcus hirae</i> | NCIMB 8192 |

RESULT

The product under test: "Surface Sanitizer Formula 80%" demonstrated bactericidal activity according to EN 1276:2019 (≥ 5 log reduction), under clean conditions for 30 seconds contact time, at 20 ± 1 °C, when tested at product concentrations:

Undiluted (80%) using as test organisms the reference strains: *Pseudomonas aeruginosa*, *Staphylococcus aureus*, *Escherichia coli* and *Enterococcus hirae*.

Results refer to the sample as received and analyzed on the period specified above.

The test report shall not be reproduced except in full, without written approval of the laboratory.

The samples will be stored by the laboratory during 1 month from the end test date.

The study report and raw data will be stored by the laboratory for 5 years.

End of Test Report